

Chiral Dynamics of Deeply Bound Pionic Atoms

E.E. Kolomeitsev,¹ N. Kaiser,² and W. Weise^{1,2}

¹*ECT*, Villa Tambosi, I-38050 Villazzano (Trento), Italy*

²*Physik-Department, Technische Universität München, D-85747 Garching, Germany*

We present and discuss a systematic calculation, based on two-loop chiral perturbation theory, of the pion-nuclear s-wave optical potential. A proper treatment of the explicit energy dependence of the off-shell pion self-energy together with (electromagnetic) gauge invariance of the Klein-Gordon equation turns out to be crucial. Accurate data for the binding energies and widths of the $1s$ and $2p$ levels in pionic ^{205}Pb and ^{207}Pb are well reproduced, and the notorious "missing repulsion" in the pion-nuclear s-wave optical potential is accounted for. The connection with the in-medium change of the pion decay constant is clarified.

PACS numbers: 36.10Gv

Keywords: pionic atoms, nuclear matter

Recent accurate data on $1s$ and $2p$ states of negatively charged pions bound to Pb nuclei [1] have set new standards and constraints for the detailed analysis of s-wave pion-nucleus interactions. This subject has a long history [2, 3, 4, 5, 6, 7, 8], culminating in various attempts to understand the notorious "missing repulsion": the standard ansatz for the s-wave pion-nucleus optical potential in terms of the empirical threshold πN -amplitudes times the proton and neutron densities $\rho_{p,n}$, supplemented by sizeable double-scattering corrections, still misses the empirically required repulsive interaction by a large amount. On purely phenomenological grounds, this problem can be fixed by simply introducing a sufficiently large negative real part, $\text{Re}B_0$, in the ρ^2 term of the pion-nuclear optical potential [6]. The arbitrariness of this procedure is of course unsatisfactory, also in view of the fact that $\text{Re}B_0$ cannot be calculated microscopically with sufficient accuracy, and that the high precision measurements of the real part of the pion deuteron scattering length [9] suggests a very small net $\text{Re}B_0$ by analogy.

In the meantime, this issue has been revived in a variety of ways. An interpretation of the "missing repulsion" in terms of a possible in-medium change of the pion decay constant, as suggested in ref. [10], appears to be remarkably successful [11, 12] but has been debated [13, 14]. Clearly, this concept needs further justification. In the present paper we show that a key to the understanding of low-energy pion-nucleus s-wave interactions lies in its distinct energy dependence imposed by chiral symmetry, in combination with the "accidental" approximate vanishing of the isospin-even threshold πN -amplitude. Another important feature, generally ignored in previous analyses, is the systematic incorporation of gauge invariance at all places where the pion energy ω appears explicitly, when solving the Klein-Gordon equation in the presence of electromagnetic interactions.

Our framework will be in-medium chiral perturbation theory at two-loop order [16]. But before going into technical details, the following simplified treatment may be useful to illustrate driving mechanisms.

Consider a zero-momentum π^- interacting with nuclear matter at low proton and neutron densities, ρ_p and ρ_n . The in-medium pion polarization operator to leading order in the nucleon densities is expressed in terms of the isospin-even and isospin-odd off-shell πN amplitudes $T^\pm(\omega)$ as $\Pi(\omega) = -T^-(\omega)\delta\rho - T^+(\omega)\rho$, with the isoscalar and isovector nucleon densities, $\rho = \rho_p + \rho_n$ and $\delta\rho = \rho_p - \rho_n$. The spontaneous and explicit breaking of chiral symmetry implies the following leading terms of those amplitudes [15]:

$$T^+(\omega) = \frac{\sigma_N - \beta\omega^2}{f_\pi^2}, \quad T^-(\omega) = \frac{\omega}{2f_\pi^2}, \quad (1)$$

where $f_\pi = 92.4$ MeV is the pion decay constant, $m_\pi = 139.57$ MeV is the (charged) pion mass and $\sigma_N \simeq 45 \pm 8$ MeV [17] is the pion-nucleon sigma-term. The empirical observation that $T^+(m_\pi) = (-0.04 \pm 0.09)$ fm $\simeq 0$ [18] sets the constraint $\beta \simeq \sigma_N/m_\pi^2$.

Next, consider the pionic in-medium dispersion equation at zero momentum, $\omega^2 - m_\pi^2 - \Pi(\omega) = 0$. Introduce an (energy independent) equivalent optical potential U by $\omega^2 = m_\pi^2 + 2m_\pi U$ with $U \ll m_\pi$, and expand around $\omega = m_\pi$. This gives

$$U \simeq \frac{\Pi(m_\pi)}{2m_\pi} \left[1 + \frac{\Pi'(m_\pi)}{2m_\pi} \right] + \dots \simeq \frac{\Pi(m_\pi)/(2m_\pi)}{\left(1 - \frac{\partial\Pi}{\partial\omega^2}\right)|_{\omega=m_\pi}}, \quad (2)$$

where the last approximate step introduces the wave function renormalization factor $\left(1 - \frac{\partial\Pi}{\partial\omega^2}\right)^{-1}_{\omega=m_\pi}$. The difference between second and last step in (2) is of sub-leading order. Inserting (1) and assuming $\delta\rho \ll \rho$ one finds

$$U \simeq -\frac{\delta\rho}{4f_\pi^2} \left(1 - \frac{\sigma_N\rho}{m_\pi^2 f_\pi^2} \right)^{-1} \simeq -\frac{\delta\rho}{4f_\pi^{*2}}, \quad (3)$$

which is the expression proposed in ref. [10]. It involves the driving Weinberg-Tomozawa term in $T^-(\omega)$, but with the pion decay constant renormalized ($f_\pi \rightarrow f_\pi^*(\rho)$) to leading order in the density ρ , in accordance with the




FIG. 1: Binding energies and widths of deeply bound pionic states in the isotopes ^{207}Pb (left figure) and ^{205}Pb (right figure). Diamonds show the experimental data from [1]. Uncertainties in the extraction of the 1s level for ^{207}Pb are indicated by different choices of a control parameter R as specified in [1]. The results for the polarization operator (8,9) are depicted by open and filled circles, respectively. Triangles show the results obtained with the chiral polarization operator (10).

The isospin-even off-shell πN -scattering amplitude at zero pion momentum can be written in the following form (for $\omega > m_\pi$):

$$T^+(\omega) = \frac{\sigma_N - \beta\omega^2}{f_\pi^2} + \frac{3g_A^2 m_\pi^3}{16\pi f_\pi^4} + \frac{3g_A^2 Q^2 m_\pi \zeta}{64\pi f_\pi^4} + iT_{\text{im}}$$

where $\beta = g_A^2/4M - 2c_2 - 2c_3$, $\sigma_N = -4c_1 m_\pi^2 - 9g_A^2 m_\pi^3/64\pi f_\pi^2$ and $T_{\text{im}} = \omega^2 Q/(8\pi f_\pi^4)$. The nucleon axial-vector coupling constant has the value $g_A = 1.27$. We introduce the abbreviation $Q = \sqrt{\omega^2 - m_\pi^2}$. The second-order low-energy constants $c_{1,2,3}$ (for notations see ref. [21]) are tuned to the empirical values of the sigma-term [17], $\sigma_N = 45$ MeV, and the πN scattering length, $T^+(m_\pi) = 0$.

The parameter ζ reflects freedom in the choice of the interpolating pion field in the effective chiral Lagrangian [13, 22]. It enters all interaction vertices with three and more pions. The one-loop correction to the (off-shell) pion self-energy in vacuum depends also on this parameter ζ . By requiring that the residue at the pion pole remains equal to one [24] as it is implicit in the form of the KG equation (4), one gets the constraint $\zeta = 0$.

The isospin-odd off-shell πN -amplitude at zero momentum reads:

$$T^-(\omega) = \frac{\omega}{2f_\pi^2} \left[1 + \frac{\gamma\omega^2}{(2\pi f_\pi)^2} \right] - \frac{\omega^2 Q}{8\pi^2 f_\pi^4} \ln \frac{\omega + Q}{m_\pi} + \frac{i}{2} T_{\text{im}} \quad (12)$$

with $\gamma = (g_A \pi f_\pi / M)^2 + \ln(2\Lambda/m_\pi)$. The cut-off scale $\Lambda = 737$ MeV $\simeq 8f_\pi$ is chosen to reproduce the central empirical value of the on-shell scattering amplitude at threshold $T^-(m_\pi) = 1.85 \pm 0.09$ fm [18]. We neglect here small additional counter-term contributions proportional to the third order low-energy constants $\bar{d}_j$ of ref. [23].

The next term in (10) corresponds to the important Ericson-Ericson double scattering correction [2] generalized to isospin asymmetric nuclear matter and off-shell pions:

$$\Pi_{\text{ds}}(\omega) = \text{---} \text{---} \text{---} \text{---} = \frac{\omega^2}{3(4\pi f_\pi)^4} \left\{ L(k_p, k_p, Q) + L(k_n, k_n, Q) + 2L(k_p, k_n, Q) \right\}, \quad (13)$$

where $k_{p,n} = (3\pi^2 \rho_{p,n})^{1/3}$ refer to the proton and neutron Fermi momenta and

$$L(k_p, k_n, Q) = 4k_p k_n (Q^2 + 3k_p^2 + 3k_n^2) + 8Q(k_n^3 - k_p^3) \ln \frac{Q + k_n - k_p}{Q - k_n + k_p} - 8Q(k_p^3 + k_n^3) \ln \frac{Q + k_p + k_n}{Q - k_p - k_n} + \left[3(k_p^2 - k_n^2)^2 + 6Q^2(k_p^2 + k_n^2) - Q^4 \right] \times \ln \frac{(k_n - k_p)^2 - Q^2}{(k_p + k_n)^2 - Q^2}. \quad (14)$$

The third term in (10) is a small relativistic correction from the particle-hole (Born) diagram evaluated at zero pion momentum:

$$\Pi_{\text{rel}}(\omega) = \text{---} \text{---} \text{---} \text{---} = \frac{g_A^2 \omega}{10(\pi M f_\pi)^2} (k_p^5 - k_n^5). \quad (15)$$

The last term in (11) represents the effect induced by $\pi\pi$ -interactions with two virtual pions being absorbed on the nucleons in the Fermi-sea, and by an additional two-loop correction [16, 22]:

$$\Pi_{\text{cor}}(\omega) = \text{---} \text{---} \text{---} \text{---} + \text{---} \text{---} \text{---} \text{---} = \frac{g_A^2}{20(4\pi f_\pi)^4}$$




FIG. 2: Dependence of energies and widths of pionic levels for ^{205}Pb on parameters of the optical potential. Solid lines correspond to the variation of the neutron radius in the interval $\delta R_n = -0.2 \div 0.2$ fm. Dashed lines show the variation with the pion nucleon sigma term $\sigma_N(0) = 20 \div 65$ MeV. Dotted lines represent the variation of $\text{Re}B_0 = (-0.5 \div 0.5) \text{Im}B_0$. Arrows indicate the directions in which results move as the varied parameters increase. Triangles and experimental points are the same as in Fig. 1.

$$\times \{ (Q^2(\zeta + 2) + 5m_\pi^2)[H(k_p, k_p) + H(k_n, k_n)] + Q^2(8\zeta - 4)H(k_p, k_n) \}. \quad (16)$$

The function $H(k_p, k_n)$ consists of the last four terms written in eq. (12) of ref. [16]. In the actual calculation the contribution $\Pi_{\text{cor}}(\omega)$ turns out to be negligibly small.

Solutions of the wave equation (4) for $1s$ and $2p$ levels using (10) are shown in Fig. 1 by triangles. They agree well with experimental data. We also find that the energy dependent polarization operator (6,7) gives equally good results as (10) if the amplitude $T^-(\omega)$ in (1) is extended to include the ω^3 term in (12), with the parameter γ tuned to the empirical value of the scattering length.

In Fig. 2 we examine the dependence of our results for ^{205}Pb on the less constrained parameters of the model. Variations of the neutron radius R_n affect mainly the binding energy, whereas the sigma term σ_N and $\text{Re}B_0$ have a stronger impact on the level width. Note the strong correlation in the effects induced by changes of $\text{Re}B_0$ and σ_N .

In summary, we have demonstrated that the long standing issue of the "missing repulsion" in the s-wave pion nucleus potential can be naturally resolved by taking into account the explicit energy dependence of the pion self-energy and the gauge invariant incorporation of electromagnetic interactions. The experimental data for $1s$ and $2p$ levels in Pb isotopes are well reproduced. We have also clarified that, to leading order, the energy dependence effects can be interpreted in terms of the in-

medium reduction of the pion decay constant, or equivalently, the renormalization of the intrinsic pion wave function in matter.

We thank E. Friedman, P. Kienle, M. Lutz, E. Oset, D. Voskresensky and T. Yamazaki for discussions and communications. We also gratefully acknowledge valuable contributions by R. Leisibach at the early stages of this work. This work is supported in part by BMBF and GSI.

-
- [1] H. Gilg et al., Phys. Rev. C62, 025201 (2000); K. Itahashi et al., Phys. Rev. C62, 025202 (2000); H. Geissel et al., Phys. Rev. Lett. 88, 122301 (2002).
 - [2] M. Ericson and T.E.O. Ericson, Ann. Phys. (N.Y.) 36, 323 (1966).
 - [3] L. Tauscher, Proc. Int Seminar on π -Meson-Nucleus Interactions, Strasbourg, 1971, 45 (CNRS, Strasbourg, 1971).
 - [4] T.E.O. Ericson and W. Weise, Pions and Nuclei (Clarendon Press, Oxford, 1988).
 - [5] A.B. Migdal, E.E. Saperstein, M.A. Troitsky and D.N. Voskresensky, Phys. Reports 192, 179 (1990).
 - [6] J. Nieves, E. Oset and C. Garcia-Recio, Nucl. Phys. A554, 509 (1993).
 - [7] T. Waas, R. Brockmann, and W. Weise, Phys. Lett. B405, 215 (1997).
 - [8] C.J. Batty, E. Friedman and A. Gal, Phys. Reports 287, 385 (1997).
 - [9] D. Chatellard et al., Nucl. Phys. A625, 855 (1997).
 - [10] W. Weise, Nucl. Phys. A690, 98 (2001); Acta Phys. Polonica B31, 2715 (2000).
 - [11] P. Kienle and T. Yamazaki, Phys. Lett. B514 (2001) 1.
 - [12] E. Friedman, Phys. Lett. B524, 87 (2002); nucl-th/0205054.
 - [13] A. Wirzba, J. Oller and U.-G. Meißner, Ann. Phys. (N.Y.) 297, 27 (2002).
 - [14] C. Garcia-Recio, J. Nieves and E. Oset, nucl-th/0202038.
 - [15] T.E.O. Ericson, Phys. Lett. B195, 116 (1987); *ibid* B321, 312 (1994).
 - [16] N. Kaiser and W. Weise, Phys. Lett. B512, 283 (2001).
 - [17] J. Gasser, H. Leutwyler and M.E. Sainio, Phys. Lett. B253, 252 (1991).
 - [18] H.Ch. Schröder et al., Phys. Lett. B 469, 25 (1999).
 - [19] G. Fricke et al., Atomic Data and Nuclear Data Tables 60, 177 (1995).
 - [20] I. Angeli, M. Beiner, R.J. Lombard and D. Mas, J. Phys. G6, 303 (1980).
 - [21] V. Bernard, N. Kaiser and U.-G. Meißner, Phys. Lett. B309, 421 (1993); Phys. Rev. C52, 2185 (1995).
 - [22] T.S. Park, H. Jung and D.P. Min, nucl-th/0101064.
 - [23] N. Fettes, U.-G. Meißner and S. Steininger, Nucl. Phys. A460 199 (1998).
 - [24] We discard here the fact the wave function renormalization factor of the pion in vacuum receives also the contribution $-2l_4 m_\pi^2 / f_\pi^2$ from the low-energy constant l_4 .